

# Neutrosophic Fuzzy Ideals in Boolean like semi rings

**.Dr.ECCLESTON**

<sup>1</sup>M. Phil Scholar, <sup>2</sup>Assistant Professor, <sup>3</sup>Associate Professor  
<sup>1,2,3</sup> PG & Research Department of Mathematics,  
A.P.C Mahalaxmi College for women, Thoothukudi.

## Abstract

The aim of this paper is to introduce the concept of Neutrosophic Fuzzy Ideals in Boolean like semi rings and also investigate some of the theorems in detail. We also obtain some characterisations and proved theorems for Boolean like semi-rings.

## Keywords:

Neutrosophic fuzzy set, Neutrosophic Fuzzy Ideal and Boolean like semi ring

## 1. Introduction

Zadeh proposed the notion of a fuzzy set in 1965. Boolean like semi rings were introduced by K. Venkateswarlu, B.V.N. Murthy and N. Amarnath. The concept of neutrosophy was introduced by Florentin Smarandache as a new branch of philosophy. Neutrosophy is a base of Neutrosophic logic which is an extension of fuzzy logic in which indeterminacy is included. In Neutrosophic logic, each proposition is estimated to have the percentage of truth in a subset T, percentage of indeterminacy in a subset I, and the percentage of falsity in a subset F. The theory of neutrosophic set have achieved great success in various fields like medical diagnosis, image processing, decision making problem ,robotics and so on. I.Arockiarani consider the neutrosophic set with value from the subset of  $[0,1]$  and extended the research in fuzzy neutrosophic set. J.Martina Jency and I.Arockia rani initiate the concept of subgroupoids in fuzzy neutrosophic set. A. Solairaju and S. Thiruvani have introduced the concept of Neutrosophic Fuzzy Ideals in near rings. R.Rajeswari and N. Meenakumari have introduced the concept of Fuzzy Bi-ideals in Boolean like semi-rings. In this paper, we recreate the concept of Neutrosophic Fuzzy Ideals in Near rings into Neutrosophic Fuzzy Ideals in Boolean like semi rings and we have discussed some of the theorems in detail.

## 2. Preliminaries

### Definition : 2.1

A non empty set  $R$  with two binary operations '+' and '·' is called a **near-ring** if

- i)  $(R, +)$  is a group
- ii)  $(R, \cdot)$  is a semigroup
- iii)  $x \cdot (y + z) = x \cdot y + x \cdot z$  for all  $x, y, z \in R$

### Definition : 2.2

A system  $(R, +, \cdot)$  a **Boolean semi ring** if and only if the following properties hold

- i)  $(R, +)$  is an additive(abelian)group(whose 'zero' will be denoted by '0')
- ii)  $(R, \cdot)$  is a semigroup of idempotents in the sense  $aa = a$ , for all  $a \in R$
- iii)  $(+)$  is commutative &
- iv)  $abc = bac$  for all  $a, b, c \in R$

### Definition : 2.3

A non-empty set  $R$  together with two binary operations  $+$  and  $\cdot$  satisfying the following conditions is called a **Boolean like semi-ring**

- i)  $(R, +)$  is an abelian group
- ii)  $(R, \cdot)$  is a semigroup
- iii)  $a \cdot (b + c) = a \cdot b + a \cdot c$  for all  $a, b, c \in R$
- iv)  $a + a = 0$ , for all  $a \in R$
- v)  $ab(a + b + ab) = ab$ , for all  $a, b \in R$

### Definition : 2.4

A non-empty subset  $I$  of  $R$  is said to be an **ideal** if,

- i)  $(I, +)$  is a subgroup of  $(R, +)$ , (i.e) for  $a, b \in R \Rightarrow a + b \in R$
- ii)  $ra \in R$  for all  $a \in I, r \in R$ , (i.e),  $RI \subseteq I$
- iii)  $(r + a)s + rs \in I$  for all  $r, s \in R, a \in I$ .

### Definition : 2.5

Let  $\mu$  be a fuzzy set defined on  $R$ . Then  $\mu$  is said to be a **fuzzy ideal** of  $R$  if

$$i) \mu(x - y) \geq \min\{\mu(x), \mu(y)\}, \text{ for all } x, y \in R$$

$$ii) \mu(ra) \geq \mu(a), \text{ for all } r, a \in R$$

$$iii) \mu((r + a)s + rs) \geq \mu(a), \text{ for all } r, a, s \in R$$

**Definition: 2.6**

A Neutrosophic fuzzy set  $A$  on the universe of discourse  $X$  characterized by a truth membership function  $T_A(x)$ , an indeterminacy function  $I_A(x)$  and a falsity membership function  $F_A(x)$  is defined as  $A = \{ \langle x, T_A(x), I_A(x), F_A(x) \rangle : x \in X \}$  where  $T_A, I_A, F_A: X \rightarrow [0, 1]$  and  $0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3$ .

**Definition: 2.7**

Let  $A$  and  $B$  be Neutrosophic fuzzy sets of  $X$ . Then

$$i) \quad A \cup B = \{ \langle x, T_{A \cup B}(x), I_{A \cup B}(x), F_{A \cup B}(x) \rangle : x \in X \}, \text{ where}$$

$$T_{A \cup B}(x) = \max\{T_A(x), T_B(x)\}, I_{A \cup B}(x) = \min\{I_A(x), I_B(x)\}, F_{A \cup B}(x) = \min\{F_A(x), F_B(x)\},$$

for all  $x \in X$ .

$$ii) \quad A \cap B = \{ \langle x, T_{A \cap B}(x), I_{A \cap B}(x), F_{A \cap B}(x) \rangle : x \in X \}, \text{ where}$$

$$T_{A \cap B}(x) = \min\{T_A(x), T_B(x)\}, I_{A \cap B}(x) = \max\{I_A(x), I_B(x)\}, F_{A \cap B}(x) = \max\{F_A(x), F_B(x)\},$$

for all  $x \in X$ .

**Definition: 2.8**

A Neutrosophic fuzzy set  $A$  in a near ring  $N$  is called **neutrosophic fuzzy sub near ring** of  $N$  if

$$i) \quad T_A(x - y) \geq \min\{T_A(x), T_A(y)\}, I_A(x - y) \leq \max\{I_A(x), I_A(y)\},$$

$$F_A(x - y) \leq \max\{F_A(x), F_A(y)\}$$

$$ii) \quad T_A(xy) \geq \min\{T_A(x), T_A(y)\}, I_A(xy) \leq \max\{I_A(x), I_A(y)\},$$

$$F_A(xy) \leq \max\{F_A(x), F_A(y)\}$$

### 3. Main Results

#### Definition: 3.1

Let  $R$  be a Boolean like semi ring. A Neutrosophic fuzzy set  $A$  in a Boolean like semi ring  $R$  is called Neutrosophic fuzzy ideal of  $R$  if

- (i)  $T_A(x - y) \geq \min\{T_A(x), T_A(y)\}, I_A(x - y) \leq \max\{I_A(x), I_A(y)\},$   
 $F_A(x - y) \leq \max\{F_A(x), F_A(y)\}$
- (ii)  $T_A(ra) \geq T_A(a), I_A(ra) \leq I_A(a), F_A(ra) \leq F_A(a)$
- (iii)  $T_A((r + a)s + rs) \geq T_A(a), I_A((r + a)s + rs) \leq I_A(a),$   
 $F_A((r + a)s + rs) \leq F_A(a)$

#### Theorem: 3.2

Let  $A$  and  $B$  be neutrosophic fuzzy ideals of  $R$ . If  $A \subset B$ , then  $A \cup B$  is a neutrosophic fuzzy ideal of  $R$ .

#### Proof:

Let  $A$  and  $B$  are Neutrosophic fuzzy ideals of  $R$ . Let  $x, y, z \in R$

- i) 
$$\begin{aligned} T_{A \cup B}(x - y) &= \max\{T_A(x - y), T_B(x - y)\} \\ &\geq \max\{\min\{T_A(x), T_A(y)\}, \min\{T_B(x), T_B(y)\}\} \\ &= \min\{\max\{T_A(x), T_B(x)\}, \max\{T_A(y), T_B(y)\}\} \\ &= \min\{T_{A \cup B}(x), T_{A \cup B}(y)\} \\ I_{A \cup B}(x - y) &= \min\{I_A(x - y), I_B(x - y)\} \\ &\leq \min\{\max\{I_A(x), I_A(y)\}, \max\{I_B(x), I_B(y)\}\} \\ &= \max\{\min\{I_A(x), I_B(x)\}, \min\{I_A(y), I_B(y)\}\} \\ &= \max\{I_{A \cup B}(x), I_{A \cup B}(y)\} \\ F_{A \cup B}(x - y) &= \min\{F_A(x - y), F_B(x - y)\} \\ &\leq \min\{\max\{F_A(x), F_A(y)\}, \max\{F_B(x), F_B(y)\}\} \\ &= \max\{\min\{F_A(x), F_B(x)\}, \min\{F_A(y), F_B(y)\}\} \\ &= \max\{F_{A \cup B}(x), F_{A \cup B}(y)\} \end{aligned}$$
- ii) 
$$\begin{aligned} T_{A \cup B}(ra) &= \max\{T_A(ra), T_B(ra)\} \\ &\geq \max\{T_A(a), T_B(a)\} \end{aligned}$$

$$\begin{aligned}
&= T_{A \cup B}(a) \\
I_{A \cup B}(ra) &= \min\{I_A(ra), I_B(ra)\} \\
&\leq \min\{I_A(a), I_B(a)\} \\
&= I_{A \cup B}(a) \\
F_{A \cup B}(ra) &= \min\{F_A(ra), F_B(ra)\} \\
&\leq \min\{F_A(a), F_B(a)\} \\
&= F_{A \cup B}(a) \\
\text{iii)} \quad T_{A \cup B}((r+a)s + rs) &= \max\{T_A((r+a)s + rs), T_B((r+a)s + rs)\} \\
&\geq \max\{T_A(a), T_B(a)\} = T_{A \cup B}(a) \\
I_{A \cup B}((r+a)s + rs) &= \min\{I_A((r+a)s + rs), I_B((r+a)s + rs)\} \\
&\leq \min\{I_A(a), I_B(a)\} = I_{A \cup B}(a) \\
F_{A \cup B}((r+a)s + rs) &= \min\{F_A((r+a)s + rs), F_B((r+a)s + rs)\} \\
&\leq \min\{F_A(a), F_B(a)\} = F_{A \cup B}(a)
\end{aligned}$$

Therefore,  $A \cup B$  is a Neutrosophic fuzzy ideal of  $R$ .

### Theorem: 3.3

Let  $A$  and  $B$  be neutrosophic fuzzy ideals of  $R$ . Then  $A \cap B$  is a neutrosophic fuzzy ideal of  $R$ .

### Proof:

Let  $A$  and  $B$  are Neutrosophic fuzzy ideals of  $R$ . Let  $x, y, z \in R$

$$\begin{aligned}
\text{i)} \quad T_{A \cap B}(x - y) &= \min\{T_A(x - y), T_B(x - y)\} \\
&\geq \min\{\min\{T_A(x), T_A(y)\}, \min\{T_B(x), T_B(y)\}\} \\
&= \min\{\min\{T_A(x), T_B(x)\}, \min\{T_A(y), T_B(y)\}\} \\
&= \min\{T_{A \cap B}(x), T_{A \cap B}(y)\} \\
I_{A \cap B}(x - y) &= \max\{I_A(x - y), I_B(x - y)\} \\
&\leq \max\{\max\{I_A(x), I_A(y)\}, \max\{I_B(x), I_B(y)\}\} \\
&= \max\{\max\{I_A(x), I_B(x)\}, \max\{I_A(y), I_B(y)\}\}
\end{aligned}$$

$$\begin{aligned}
&= \max\{I_{A \cap B}(x), I_{A \cap B}(y)\} \\
F_{A \cap B}(x - y) &= \max\{F_A(x - y), F_B(x - y)\} \\
&\leq \max\{\max\{F_A(x), F_A(y)\}, \max\{F_B(x), F_B(y)\}\} \\
&= \max\{\max\{ ( ), ( ) \}, \max\{ ( ), ( ) \}\} \\
&= \max\{F_{A \cap B}(x), F_{A \cap B}(y)\} \\
\text{ii)} \quad T_{A \cap B}(ra) &= \min\{T_A(ra), T_B(ra)\} \\
&\geq \min\{T_A(a), T_B(a)\} \\
&= T_{A \cap B}(a) \\
I_{A \cap B}(ra) &= \max\{I_A(ra), I_B(ra)\} \\
&\leq \max\{I_A(a), I_B(a)\} \\
&= I_{A \cap B}(a) \\
F_{A \cap B}(ra) &= \max\{F_A(ra), F_B(ra)\} \\
&\leq \max\{F_A(a), F_B(a)\} \\
&= F_{A \cap B}(a) \\
\text{iv)} \quad T_{A \cap B}((r + a)s + rs) &= \min\{T_A((r + a)s + rs), T_B((r + a)s + rs)\} \\
&\geq \min\{T_A(a), T_B(a)\} = T_{A \cap B}(a) \\
I_{A \cap B}((r + a)s + rs) &= \max\{I_A((r + a)s + rs), I_B((r + a)s + rs)\} \\
&\leq \max\{I_A(a), I_B(a)\} = I_{A \cap B}(a) \\
F_{A \cap B}((r + a)s + rs) &= \max\{F_A((r + a)s + rs), F_B((r + a)s + rs)\} \\
&\leq \max\{F_A(a), F_B(a)\} = F_{A \cap B}(a)
\end{aligned}$$

Therefore,  $A \cap B$  is a Neutrosophic fuzzy ideal of  $R$ .

#### Lemma: 3.4

For all  $a, b \in I$  and  $i$  is any positive integer, if  $a = b$ , then

- i)  $a^i \leq b^i$
- ii)  $[\min( , )] = \min( , )$
- iii)  $[\max( , )] = \max( , )$

**Theorem: 3.5**

Let  $A$  be a Neutrosophic fuzzy ideal of  $R$ . Then  $A^m = \{ \langle x, T_A(x), I_A(x), F_A(x) \rangle : x \in R \}$  is a Neutrosophic fuzzy ideal of  $R^m$ , where  $m$  is a positive integer and  $T_{A^m}(x) = (T_A(x))^m, I_{A^m}(x) = (I_A(x))^m, F_{A^m}(x) = (F_A(x))^m$ .

**Proof:**

Let  $A$  be a Neutrosophic fuzzy ideal of  $R$ . Let  $x, y, z \in R$ .

$$\begin{aligned}
 \text{i)} \quad T_{A^m}(x - y) &= (T_A(x - y))^m \\
 &\geq [\min\{T_A(x), T_A(y)\}]^m \\
 &= \min\{ (T_A(x))^m, (T_A(y))^m \} \\
 I_{A^m}(x - y) &= (I_A(x - y))^m \\
 &\leq [\max\{I_A(x), I_A(y)\}]^m \\
 &= \max\{ (I_A(x))^m, (I_A(y))^m \} \\
 F_{A^m}(x - y) &= (F_A(x - y))^m \\
 &\leq [\max\{F_A(x), F_A(y)\}]^m \\
 &= \max\{ (F_A(x))^m, (F_A(y))^m \}
 \end{aligned}$$

$$\begin{aligned}
 \text{ii)} \quad (T_A(a))^m &= (T_A(a))^m \\
 &\geq (T_A(a))^m \\
 &= (T_A(a))^m \\
 (I_A(a))^m &= (I_A(a))^m \\
 &\leq (I_A(a))^m \\
 &= (I_A(a))^m \\
 (F_A(a))^m &= (F_A(a))^m \\
 &\leq (F_A(a))^m \\
 &= (F_A(a))^m
 \end{aligned}$$

$$\begin{aligned}
 \text{iii)} \quad ((T_A(a) + T_A(b))^m) &= ((T_A(a) + T_A(b))^m) \\
 &\geq (T_A(a))^m \\
 &= (T_A(a))^m \\
 ((I_A(a) + I_A(b))^m) &= ((I_A(a) + I_A(b))^m)
 \end{aligned}$$

$$\begin{aligned}
&\leq (I_A(a)) \\
&= ( ) \\
(( + ) + ) &= ( (( + ) + )) \\
&\leq (F_A(a)) \\
&= ( )
\end{aligned}$$

Therefore, is a Neutrosophic fuzzy ideal of .

#### 4. References

- [1] S.Abu zaid, On fuzzy subnear-rings and ideals, Fuzzy sets and systems 44(1991)139-146
- [2] Agboola A.A.A, Akwu A.D and Oyebo Y.T., Neutrosophic groups and subgroups, International J.Math.Combin.Vol.3(2012),1-9.
- [3] I.Arockiarani, J.Martina Jency , More on Fuzzy Neutrosophic sets and Fuzzy Neutrosophic Topological spaces , International journal of innovative research and studies,May(2014), vol 3,Issue 5, 643-652.
- [4] V.Chinnadurai and S.Kadalarasi,Direct product of fuzzy ideals of near-rings,Annals of Fuzzy Mathematics and Informatics,2016
- [5] T.K.Dutta and B.K.Biswas, Fuzzy ideal of a near-ring,Bull.Cal.Math.Soc.89(1997),447-456.
- [6] S.D.Kim and H.S.Kim , On fuzzy ideals of near-rings, Bulletin Korean Mathematical society 33(1996) 593-601
- [7] J.Martina Jency , I.Arockiarani, Fuzzy Neutrosophic Subgroupoids,Asian Journal of Applied Sciences(ISSN:2321-0893),vol 04,Issue 01,February (2016)
- [8] R.Rajeswari, N.Meenakumari, *Fuzzy ideals in Boolean like Semi-rings* in Enrich, Vol.V (II), 60-66(2014).
- [9] Rosenfield A, Fuzzy Groups,Journal of mathematical analysis and applications,35,512-517(1971)



- [10] F.Smarandache, Neutrosophy ,A new branch of Philosophy logic in multiple-valued logic,An international journal,8(3)(2002),297-384.
- [11] A. Solairaju, S. Thiruveni, Neutrosophic Fuzzy Ideals in near rings, International Journal of Pure and Applied Mathematics, Volume 118 No.6 2018, 527-539
- [12 ] K.Venkateswarlu and B.V.N.Murthy and Amaranth, *Boolean like semi rings*, Int.J.Contemp.Math.Sciences, Vol.6,2011,no.13,619-635.
- [13] Zadeh.L.A, Fuzzy sets, Information control, Vol.8,338-353(1965)