

## ANTIMICROBIAL ACTIVITY OF LACTIC ACID BACTERIA AGAINST MULTIPLE DRUG RESISTANT UROPATHOGENS

**Dr.ECCLESTON**

<sup>1</sup>\*Research scholar, Microbiology Research Laboratory,  
Rajasthan Aryans Mahavidyalaya, Washim, Maharashtra, India

**Dr DINESH KUMAR**

<sup>2</sup>Associate professor and Head, U.G., PG. and Research section,  
Department of Microbiology,  
Rajasthan Aryans Mahavidyalaya, Washim, Maharashtra, India

### ABSTRACT

Urinary tract infections (UTIs) are the common bacterial infections that affect various areas of the urinary system in both females and males. Globally, the morbidity rates associated with UTI are very high due to occurrence of antimicrobial resistance in pathogens causing UTI. Many studies have been conducted to overcome the resistant in UTI pathogens. One of the alternatives for treating UTI pathogens are Lactic acid bacteria (LAB) commonly known as probiotics. The present research aims to explore the bio therapeutic application of LAB over Uropathogens in order to reduce the overreliance on antibiotics. In the present study, LAB was isolated from different dairy products viz. buttermilk, yoghurt and curd. Four different LAB isolates were identified as *Lactobacillus plantarum*, *Lactobacillus rhamnosus*, *Lactobacillus casei* and *Lactococcus lactis*. The results revealed that maximum antimicrobial activity was given by *Lactobacillus plantarum* and *Lactobacillus rhamnosus* against most of the Uropathogens. Hence, Probiotics could be utilized as one of the best natural therapeutics for prevention as well as cure of urinary tract infections.

**Key words:** UTI, LAB, Antimicrobial resistance, Probiotics, Uropathogens

### I. INTRODUCTION

Urinary tract infections are the frequent infections affecting most of the population. It mostly includes Cystitis, Pyelonephritis and Urethritis. The infection is mostly caused by one or more microorganisms belonging to aerobic or facultative bacteria, yeast and molds or sometimes viruses also. Most of the studies had reported that 80 to 85% UTI is caused due to *E.coli*. Similarly other Uropathogens reported are *Candida spp.*, *Staphylococcus spp.*, *Klebsiella spp.*, *Enterococci spp.*, and *Proteus mirabilis*. [1-3]

Females are more vulnerable to UTI as compare to males due to urogenital tract anatomy. It is reported that every year more than 40% women develop UTI. [4] The recurrence of UTI has also been reported. However, the mortality rates are less but morbidity is high. [5] The symptoms of UTI include frequent and urgent need of urination, burning or painful micturition, bad-smelling urine, pus or blood in urine, lower abdominal pain, pressure or cramps. If the infection spreads to the kidneys, symptoms could include fever, chills, nausea, vomiting, and fatigue in addition to mid-back pain (to the right or left of the spine). The risk group includes sexually active young women, patients undergoing genitourinary instrumentation or catheterization and several other populations. Typically ampicillin or a combination of trimethoprim and Sulfamethoxazole [TMP/SMX] are the usual treatments for this infection. However, development of antibiotic resistance in pathogens associated with

UTI makes the antibiotic therapy failure. Hence, search for new therapeutic agent is needed.

Previous studies had reported that lactic acid bacteria (LAB) known for their probiotic activity has potential to inhibit Uropathogens.[6-7] They act as a shield against infection forming a natural barrier in urinary system and can prevent the occurrence of UTI. Numerous antibacterial properties are displayed by lactic acid bacteria due to production of organic acids, and other compounds such as bacteriocins and antifungal peptides. [8-10] So far there is little information on the LAB isolated from dairy products against organisms implicated in UTI. Hence, the present study aims to explore the potential of LAB to treat Uropathogens to decrease the over reliance on antimicrobials.

## II. MATERIAL AND METHOD

### 2.1. Isolation of Uropathogens -

The urine samples of UTI positive patients were collected from different pathology laboratories located at Washim city area and immediately transported to Microbiology laboratory, R. A. College Washim Under aseptic condition. The urine samples were enriched in sterile nutrient Broth in aseptic condition. Then the broth was allowed to incubate at 37°C for 24 hours. After 24 hours, the turbidity was observed for the enrichment of Uropathogens.

The enriched sample was spread on different Selective media plates viz. EMB, MSA, MacConkey, Cetrimide and Potato dextrose agar plates for the isolation of Uropathogens. After incubation, uropathogens were identified by adopting conventional identification methods viz. Staining techniques (Gram staining and Lactophenol cotton blue staining), motility, colony characters Biochemical tests, IMViC, Sugar fermentation tests, enzyme assay etc. and were confirmed by comparing with Bergey's manual of systematic bacteriology.

### 2.2. Antibiotic Susceptibility Test of Uropathogens -

The antibiotic susceptibility of uropathogens was determined by Kirby Bauer disc diffusion technique suggested by CLSI (Clinical Laboratory Standards Institute). The uropathogens were separately inoculated in sterile nutrient broth tubes and incubated at 37°C for 3 to 4 hours and the turbidity was compared with 0.5 McFarland standards. The uropathogens were spread on the surface of sterile Muller- Hinton Agar plates. The antibiotic discs namely Tetracycline (30µg), Azithromycin (15µg), Cephalexin (30µg), penicillin (10U) were placed aseptically on the surface of seeded agar plates and incubated at 37°C for 18 to 24 hours. After incubation the plates were observed for the zone of inhibition and the diameter of zones was measured using HiAntibiotic Zone Scale C. [11]

### 2.3. Isolation and identification of Lactic Acid Bacteria from Dairy products-

The sealed dairy samples viz. Yoghurt, Curd and Buttermilk was purchased from Local Dairy located in Washim city area and transported to R. A. College, Microbiology Laboratory.

1ml Yoghurt and Curd sample were separately diluted in 9 ml. sterile distilled water by serial dilution. 0.1 ml. each dairy samples were spread on MRS agar plates. These Petri plates were then incubated at 37°C for 48 hrs. in bacteriological incubator. After incubation the diversified colonies grown over the MRS media were streaked on MRS agar plates. The isolated and purified cultures were subjected for conventional identification using different tests viz. Gram staining, motility, colony characters, biochemical and sugar fermentation pattern, etc. The bacterial cultures were confirmed by comparing the characters with Bergey's manual of bacteriology.

#### 2.4. Antimicrobial activity of LAB against Uropathogens-

The antimicrobial activity of LAB against uropathogens was determined by Kirby Bauer disc diffusion technique suggested by CLSI (Clinical Laboratory Standards Institute) as describe earlier. [12]

### III. RESULT AND DISSCUSION

The purpose of this study was to evaluate the antimicrobial activity of Lactic Acid Bacteria on the MDR pathogens associated with Urinary tract infection. The uropathogens were isolated from urine samples and identified as *E.coli*, *Proteus species*, *Klebsiella species*, *Staphylococcus species*, *Pseudomonas species* and *Candida species* respectively. The antibiotic susceptibility of these Uropathogens was analysed.

Table.1 and fig.1 represents the findings on the antibiotic susceptibility pattern of isolated Uropathogens. From the table it is observed that all the Uropathogens showed resistance against the tested antibiotics. In tetracycline, the minimum zone diameter (7 mm) was exhibited by *Pseudomonas species* and maximum zone of inhibition (11 mm) was shown by *Staphylococcus species* and *Candida spp.* respectively. The mean zone of inhibition against tetracycline was found to be 9.83 mm.


In case of Penicillin, the zone of inhibition (9 mm) was at par for *Pseudomonas species*, *Proteus species* and *Klebsiella species* respectively and maximum zone of inhibition was 11 mm shown by *Staphylococcus spp.* and *Candida spp.* Hence, the mean zone diameter is 9.83 mm which denotes the resistance of uropathogens. Considering Azithromycin, the minimum diameter of zone of inhibition was 9 mm shown by *Proteus spp.* and *Klebsiella*. The maximum zone of inhibition was 15 mm shown by *Staphylococcus spp.* Hence, the mean zone diameter is 11.16 mm which denotes the resistance of uropathogens.

Similarly, in Cephalixin minimum zone was 9 mm shown by *Pseudomonas species* and *Staphylococcus species* while the maximum zone of inhibition was 11 mm shown by *E.coli*. Hence, the mean zone diameter is 9.83 mm. The maximum zone of inhibition was exhibited by *Staphylococcus species* (11.5 mm) followed by *E Coli* (11.25 mm) and *Candida spp.* (10.5 mm). *Proteus* and *Klebsiella* showed at par results of inhibition zone against all the antibiotics. *Pseudomonas spp.* exhibited least zone of inhibition (8.75 mm) depicting its high resistance towards antibiotics (Fig.2), from the above results, it is concluded that all the Uropathogens are Multidrug Resistant (MDR). The results on present studies are in accordance with the experimental findings of most of the research workers enlighten same line of research. [13-16]

**Table1:AntibioticsusceptibilitypatternofUropathogens**

Antibiotics	Tetracycline (30µg)	Penicillin (10U)	Azithromycin (15µg)	Cephalexin (30µg)	MeanZone ofinhibitio n
Microorganism	Zoneofinhibitioninmm				
<i>E.coli</i>	10(R )	10(R )	14 (I)	11 (R)	11.25
<i>Proteus spp.</i>	10(R )	9(R )	9(R)	10(R )	9.5
<i>Klebsiella spp.</i>	10(R )	9(R )	9(R)	10(R )	9.5
<i>Pseudomonas spp.</i>	7(R )	9(R )	10(R)	9(R)	8.75
<i>Staphylococcuspp.</i>	11(R )	11(R )	15 (I)	9(R)	11.5
<i>Candida spp.</i>	11(R )	11(R )	10(R)	10(R )	10.5
<b>Mean zone ofinhibit ion</b>	9.83	9.83	11.16	9.83	

{R-ResistantandI-Intermediate}

**Figure1: AntibioticsusceptibilitypatternofUropathogens**


**Figure2:Meanzoneofinhibitionexhibitedbyuropathogens**

Table 2 represents the results on identification of LAB isolated from different dairy samples. From the table it is observed that four different LAB belonging to different species were isolated from the dairy samples. The isolates were labelled according to their source of isolation as Y1 (Yoghurt), C1 (Curd), B1 & B2 (Buttermilk). From the observations obtained by Biochemical tests, catalase test, Carbohydrate Utilization test, cultural characteristics and Gram Staining processes, the Y1 isolate identified as *Lactobacillus casei*, C1 as *Lactococcus lactis*, B1 & B2 as *Lactobacillus rhamnosus* and *Lactobacillus plantarum* respectively. [17]

**Table2:-IdentificationofLABbyconventionalmethod**

Culturalcharacters	Isolatesfromdairysamples			
	Y1	C1	B1	B2
Size	1.1x4.0mm	1mm	1.0-1.5mm	0.9×3um
Shape	Convex, Shortchains	SmallCircular	Round	SmallCircular
Colour	Opaquewithoutpigment	Whitecreamy	White	White
Type	Smooth	Smooth	Smooth	Liquid
Gramnature	Grampositive	Grampositive	Grampositive	Grampositive
Shape	Rod	Cocci	Rod	Rod
<b>BiochemicalCharacters</b>				
Indole	-	-	-	-
Methylred	-	-	-	-
Voges-Proskauer	-	-	-	-
Citrate	+	+	-	-
Catalase	-	-	-	-
Glucose	+	+	-	+
Lactose	+	+	+	+
Sucrose	+	-	-	+

Possible species	<i>Lactobacillus casei</i>	<i>Lactococcus lactis</i>	<i>Lactobacillus rhamnosus</i>	<i>antarum</i>
------------------	----------------------------	---------------------------	--------------------------------	----------------



Table.3 represents the findings on antimicrobial activity of LAB against UTI Pathogens. From the table it was observed that *Lactobacillus plantarum* followed by *Lactobacillus rhamnosus* isolated from buttermilk showed more antimicrobial activity against uropathogens. The average zone of inhibition exhibited by both was found to be 12.83 mm and 12.66 mm respectively. *Lactobacillus casei* isolated from yoghurt showed average zone diameter of 12.16 mm and *Lactococcus lactis* from curd showed average zone of 10.33 mm respectively. Hence, buttermilk was found to possess more benefit as compared to curd and yoghurt. However, the variations in findings might be possible because the growth of LAB and its beneficial effects are dependent on production condition, environmental temperature, external factors, quality of raw material for product preparation etc.

The mean zone of inhibition given by *Staphylococcus spp.* against all LAB was found to be (13.75 mm) followed by *E. coli* (12.25 mm), *Klebsiella spp.* (12.5 mm), *Proteus spp.* (12 mm) and *Candida spp.* (11.25 mm). *Pseudomonas spp.* showed least zone of inhibition (10.25 mm) (Fig 4). Hence, LAB was found to be effective against uropathogens.

The results on present studies are in accordance with the experimental findings of most of the research workers enlighten same line of research. They reported that Lactobacilli can prevent the adherence, growth and colonization of uropathogenic bacteria. [18-22]

**Table3:-Antimicrobial Activity of Lactic Acid Bacteria against Uropathogens.**

LAB	<i>Lactobacillus casei</i>	<i>Lactococcus lactis</i>	<i>Lactobacillus rhamnosus</i>	<i>Lactobacillus plantarum</i>	Mean zone of Inhibition (mm)
<b>Uropathogens</b>	Zone of inhibition in mm.				
<i>E. coli</i>	13	10	14	12	12.25
<i>Proteus Spp.</i>	14	10	11	13	12
<i>Klebsiella</i>	12	10	13	15	12.5
<i>Pseudomonas spp.</i>	9	12	11	9	10.25
<i>Candida spp.</i>	10	11	11	13	11.25
<i>Staphylococcus spp.</i>	15	9	16	15	13.75
<b>Average Zone (mm)</b>	12.16	10.33	12.66	12.83	

**FIG3: Antimicrobial Activity of Lactic Acid Bacteria against Uropathogens****FIG4: Mean Antimicrobial activity of LAB against Uropathogens.**

#### IV. CONCLUSION

The present study concluded that the species of Lactic acid bacteria isolated from different dairy products viz. Yoghurt, Curd and Buttermilk were found to be a perfect substitute to antibiotics used for the treatment of multiple drug resistant UTI pathogens such as *Candida spp.*, *Klebsiella spp.*, *E.coli*, *Pseudomonas spp.* and *Staphylococcus Spp.* Hence, Lactic acid bacteria can prove to be useful if consumed as Probiotics against Urinary tract infections.

## V. REFERENCES

- [1] Vasudevan R., "Urinary tract infection: an overview of the infection and the associated risk factors." *J Microbiol Exp.* 2014; 1(2):42-54. DOI:10.15406/jmen.2014.01.00008
- [2] Stamm, W.E. and T.M. Hooton, "Management of urinary tract infection in adults." *New Engl. J. Med.*, 329:1328-1334, 1993.
- [3] Jenny Stamos Kovacs, *Urinary Tract Infections (UTIs)* -webMD
- [4] Kunin, C.M., "Urinary tract infections in females." *Clin. Infect. Dis.*, 18:1-12, 1994.
- [5] Betsy Foxman, "Epidemiology of urinary tract infections: incidence, morbidity, and economic costs." *The American journal of medicine* 113(1), 5-13, 2002.
- [6] Sanni, A.I., A.A. Onilude, S.T. Ogunbanwo and S.F. Smith, "Antagonistic activity of bacteriocin produced by *Lactobacillus* species from Ogi, an indigenous fermented food." *J. Basic Microbiol.*, 39:189-195, 1999.
- [7] Reid, G., D. Buerman, C. Heinemann and A.W. Bruce, "Probiotic *Lactobacillus* dose required to restore and maintain a normal vagina flora." *FEMS Immunol. Med. Microbiol.*, 32:37-41., 2001.
- [8] Luc De Vuyst, Frédéric Leroy, "Bacteriocins from lactic acid bacteria: production, purification, and food applications." *J Mol Microbiol Biotechnol.* 2007.
- [9] Nicolle, L.E., "Asymptomatic bacteriocin - important or not?" *New Engl. J. Med.*, 343:1037-1039. 2000
- [10] Mduduzi Paul Mokoena, "Lactic Acid Bacteria and Their Bacteriocins: Classification, Biosynthesis and Applications against Uropathogens: A Mini-Review." *Molecules*; 22(8):1255. Published online 2017 Jul 26. Doi:10.3390/molecules22081255, Aug 2017.
- [11] In Seok Lim, Ho Seok Lee, and Won Yong Kim, "The Effect of Lactic Acid Bacteria Isolates on the Urinary Tract Pathogens to Infants In Vitro." *J Korean Med Sci.*; 24(Suppl 1):S57-S62, Jan 2009.
- [12] Bolanle A Adeniyi, Funmilola A Ayeni, Ogunbanwo S.T., "Antagonistic Activities of Lactic Acid Bacteria Isolated from Nigerian Fermented Dairy Food Against Organisms Implicated in Urinary Tract Infection." *Biotechnology (Faisalabad)* 5(2) DOI:10.3923/biotech.2006.183.188. Feb 2006.
- [13] Chandrika Dasgupta, Md. Abdur Rafi, and Md. Abdus Salam, "High prevalence of multidrug resistant uropathogens: A recent audit of antimicrobial susceptibility testing from a tertiary care hospital in Bangladesh." *Pak J Med Sci.* 2020 Sep-Oct; 36(6):1297-1302.
- [14] Slack, R.C.B., "Definition of urinary tract infection and assessment of efficacy in drug trials - a laboratory perspective." *Infection*, 20(Suppl 3), S155-S156., 1992
- [15] Mathai D., *et al.*, "Epidemiology and frequency of resistance among pathogens causing urinary tract infections in 1,510 hospitalized patients." *Diagn. Microbiol. Infect. Dis.* 40:129-136, 2001.
- [16] Catal., *et al.* 2008, "The increase in antimicrobial resistance shown by the uropathogens and found multiple drug resistance (MDR) in uropathogens."
- [17] Gregor Reid, Duane Charbonneau, Julie Erb, Barbara Kochanowski, Dee Beuerman, Russ Poehner, Andrew W. Bruce, "Oral use of *Lactobacillus rhamnosus* GR-1 and *L. fermentum* RC-14 significantly alters vaginal flora." *FEMS Immunology & Medical Microbiology*, Volume 35, Iss



ue2, March 2003

[18]

Asma Manzoor, Ikram Ul-

Haq, Shahjhan Baig, Javed Iqbal Qazi and Sanja Seratlic, "Efficacy of Locally Isolated Lactic Acid Bacteria against Antibiotic-Resistant Uropathogens." Jundishapur J Microbiol. 2016.

[19]

Jacqueline A. McGroarty and Gregor Reid, "Detection of a lactobacillus substance that inhibits *Escherichia coli*." Publication: Canadian Journal of Microbiology, August 1988.

[20]

Atiyeh Naderi, Roha Kasra-

Kermanshahi, Sara Gharavi, Abbas Ali Imani Fooladi, Meghdad Abdollahpour Alitappeh, Parvaneh Saffarian, "Study of antagonistic effects of Lactobacillus strains as probiotics on multidrug resistant (MDR) bacteria isolated from urinary tract infections." Iran J Basic Med Sci. 2014 Mar.

[21] Yoon Hee Shim, Seung Joo Lee, Jung Won Lee, "Antimicrobial activity of lactobacillus strains against uropathogens." Pediatr Int. 2016 Oct.

[22] Z Heshmatipour, H Ashlikousha, F Mohamaddiebli, Ashrafi A Eslami, "Antibacterial activity lactic acid bacteria (LAB) isolated native yoghurt against ESBL producing *E. coli* causing urinary tract infection." International Journal of Life Sciences Biotechnology And Pharma Research 4 (2), 117, 2015.