

Acute Toxicity of Naphthalene on a local fresh water fish – *Puntius ticto* (Ham.)

Dr DINESH KUMAR, Department of Zoology, St. Francis de Sales College, Nagpur, 440 006 Maharashtra, India.

Abstract:

Polycyclic Aromatic Hydrocarbons (PAHs) have become widespread pollutants of water originating mainly from petroleum products. These get accumulated in tissues and are converted into secondary metabolites which show bioaccumulation in the living systems and have several adverse effects on them. They further get biomagnified up through food chains. The present study attempts to evaluate the toxicity of one such PAH, Naphthalene, on the fresh water fish, *Puntius ticto*, by using pure Naphthalene dissolved in dichloromethane. LC₅₀ values were seen in the range of 3.0-3.5 after exposure of fish to Naphthalene for 24 to 96 hours. Statistical analysis revealed regression values in the range of 0.982 to 0.994, indicating high correlation between the mortality rate and concentration of Naphthalene. This paper discusses in detail the behavioral changes observed along with study of acute toxicity of Naphthalene on fresh water fish *Puntius ticto*.

Key words: PAHs; bioassay; toxicity; *Puntius ticto*.

Introduction:

In the recent decades, toxicity evaluation of industrial and domestic effluent has gained significant importance in the field of ecotoxicology. Fossil fuel consumption has increased many folds due to increased industrialization, population growth and man's greed for comfort. Oil refineries add to the woes of pollution. Due to inefficient combustion of carbonaceous matter and PAHs released from it there is ultimate effect upon the aquatic environment¹. Apart from these factors, industrial and domestic sewage effluents, surface run-off from land, deposition of airborne particulates and spillage of petroleum and petroleum products adds to the aquatic pollution. PAHs are common environmental pollutants found in automobile emissions, tobacco smoke, charcoal broiled food and chimney soot. Many of these pollutants are an important class of environmental pollutants that are known to exert carcinogenic and immunotoxic effects on aquatic organisms². PAHs as such are relatively inert compounds that are metabolized by living systems and the metabolites so formed are the actual agents which often prove to be mutagenic and carcinogenic in nature³. Thus, PAHs are ubiquitous trace elements of marine and freshwater environmental pollutants that are potent mutagens and carcinogens in nature⁴. Literature reports fish kills and also accumulation of PAHs by fish *Tilapia mossambicus*⁵. Reports are available on the effects of PAH on fish due to oil spills⁶.

In recent years, more attention has been focused on acute toxicity evaluation of different industrial waste waters. Fish bioassay is widely used to study the toxicity of different xenobiotics. PAH as a xenobiotic persists in nature for long period of time and gets bioaccumulated in aquatic biota including fish. Residues of PAH were found in aquatic species from few rivers and estuarine systems⁷. PAHs have deleterious effects on fish species⁹. Apart from impairment of water quality due to these substances, the aquatic organisms also suffer when PAH laden waste water is discharged into other water bodies.

At this juncture, it is necessary to carry out fish toxicity/ bioassays to assess the toxicity levels. Fish bioassay is widely used to study the toxicity of xenobiotics. Bioassay results provide baseline information that might be helpful in formulating strategies for dilution and controlled release of treated effluents into the recipient water bodies. Over the years, oil spills have been associated unequivocally with damage to aquatic organisms and marine birds higher up the food chain. Oil spills are the main source of PAH in the aquatic environment. Polycyclic Aromatic Hydrocarbons are produced by most

oil refineries and gas installations and significant quantities of affected water are generated. This waste water requires treatment prior to its discharge. This water, if not treated properly, leads to pollution in the receiving waters.

Report on acute toxicity of seven PAHs namely acenaphthalene, chrysene, fluoranthene, fluorine, naphthalene, phenanthrene and pyrene on benthic amphipod is available in literature, where LC₅₀ and NOEC based on 96 hrs acute toxicity have been calculated for the benthic amphipod¹⁰. However, literature on acute toxicity bioassay tests for evaluating LC₅₀ concentration of PAHs on fish is very scanty. Few studies on acute toxicity of retene to two freshwater species of Zebrafish and rainbow trout are reported.

Actually, most acute toxicity studies reported were for a longer period of 14 days, and larval stages of the fish were used¹¹. Fish bioassay to evaluate the toxicity of benzo[a] pyrene to teleost *Fundulus heteroclitus* has been quoted in literature but these tests were of long duration of 21 days and the fish were fed with pyrene contaminated *Nereis*¹². Studies were more focused on the effects due to ingestion of PAH contaminated food.

As such, literature on LC₅₀ values obtained from short term 96-hour acute toxicity tests on fish due to PAHs are rare. Majority of the work pertains to long time chronic effects of contaminated soils and water on fish. To understand the effects of exposure of fish to PAH concentrations present in waters polluted due to oil spills, detailed laboratory studies by acute toxicity tests were made to arrive at LC₅₀ values using pure PAH i.e. Naphthalene, which would give an insight into the toxicity levels of PAH. Fewer studies have been carried out in fish to determine such acute toxicity. Considering the PAH pollution due to natural disasters, it was envisaged to evaluate acute toxicity of Naphthalene using the local fresh water fish *Puntius ticto*.

Materials and Methods:

The test fish *Puntius ticto* were procured from a fresh water lake. Methods for measuring the toxicity of industrial effluents/ chemicals on fish were followed as cited in literature^{13, 14}. The fish were acclimatized for ten days in aerated and dechlorinated tap water (dilution water) at ambient temperature. Analysis of dilution water was carried out as per standard method (AHPA-1998)¹⁵ and the characteristics of dilution water are shown in Table I.

Once the fishes were acclimatized, acute toxicity tests were initiated. Controls were maintained in duplicate. During acute toxicity evaluation, the test water was changed daily and fresh dose of Naphthalene was introduced into it. Solution of Naphthalene was prepared using Dichloromethane and the same amount of dichloromethane was also added in the control.

10 fishes for each concentration were used in 10-liter aquariums and mortality rate was monitored every 24 hours for a period of 96 hours. Percent mortality was plotted against the Naphthalene concentration on probability paper and LC₅₀ values were graphically derived. Fig. 1 shows Graph indicating effect of Naphthalene on fish mortality up to 96 hours. Using Litchfield and Wilcoxon method¹⁶, 95% confidence limit, both upper and lower limits were calculated. The Slope function (S) was calculated as indicated in the literature and regression (R²) values were also calculated. Results obtained are indicated in Table II.

Results and Discussion:

The fresh water fish *Puntius ticto* is supposed to be very delicate. This was observed during the acute toxicity tests. It exhibited several types of reactions. For instance, when low Naphthalene concentrations were added, fish exhibited distress during the initial stages. They became disorientated and began to swim upside down and also exhibited somersaulting immediately on the addition of chemicals. Later, they gradually regained posture and started swimming normally. At higher dosages,

the fish initially settled at the bottom. Opercular movements became very slow and the fish were seen to very frequently come to the surface to gulp air due to respiratory distress. Additionally, bulging of eyeballs and the abdomen were very conspicuous features on exposure to high concentrations of Naphthalene.

Static bioassay tests indicate that the toxicity of Naphthalene to *Puntius ticto* is a function of the Chloride molecule present and the exposure period. Acute toxicity bioassay tests, the results of which are expressed as median lethal concentration (LC₅₀) provide the means to calculate the relative toxicity of industrial effluents of various chemicals and the relative tolerance of various fishes. The results obtained were subjected to statistical evaluation. Correlation coefficients, slope function and confidence interval were also calculated.

Table I: Characteristics of Dilution Water

Parameters	Values *
Temperature ° C	25-27
pH	7.5-8.2
Total Alkalinity as CaCO ₃	156-190
Total Hardness as CaCO ₃	142-172
Ca Hardness as CaCO ₃	80-94
Mg Hardness as CaCO ₃	62-78
Dissolved Oxygen	6.9-7.3
Calcium as Ca	32-38
Magnesium as Mg	14-18
Sodium as Na	36-38
Potassium as K	2-4
Chloride as Cl	126

(All the values are expressed as mg/L except temperature and pH.)

Regression values ranged between 0.982 to 0.994, indicating good correlation between fish mortality and PAH concentrations. Sensitivity of an organism to a toxic chemical/ effluent cannot be judged only by comparing LC₅₀ of the chemicals/ effluents. The full range of lethal concentration (LC₀-LC₁₀₀) should be taken into consideration while assessing the susceptibility of organism to any toxic chemical/ effluent. Results obtained from bioassay studies will help more scientific management by industry for taking necessary pollution control measures before the discharge of the effluents into the natural streams or water bodies. This would help in minimizing many ecotoxicological problems. Results of fish bioassay, widely used to study the toxicity of xenobiotics, provide baseline information in formulation of strategies for controlled release of treated industrial effluents into receiving water bodies. For application of toxicity data in regulation of waste water discharge and the prediction of environmental effects- both acute and chronic, toxic levels have to be determined to conserve aquatic life.

Based on per cent mortality of fish *Puntius ticto* over different periods of exposure to Naphthalene at various concentrations, slope function and regression were calculated graphically.




Figure 1. Graph showing slope function and regression values for *Puntius ticto* exposed to Naphthalene

Based on the observations, LC₅₀, NOEC, 95% confidence interval, slope function and regression were calculated for Naphthalene, and are shown in Table II.

Table II: LC₅₀, NOEC, 95% Confidence Limit, Slope function and regression values for Naphthalene to *Puntius ticto*

Exposure Period	Parameter	Values
24 hrs	LC ₅₀ mg/L	4.2
	NOEC mg/L	3.4
	95% Confidence Interval	3.44-5.124
	Slope function	$Y=12.333x-77.778$
	Regression (R ²)	0.9896
48 hrs	LC ₅₀ mg/L	3.9
	NOEC mg/L	3.0
	95% Confidence Interval	3.36-4.524
	Slope function	$Y=11.455x-46.364$
	Regression (R ²)	0.994
72 hrs	LC ₅₀ mg/L	3.5
	NOEC mg/L	2.6
	95% Confidence Interval	3.30-3.71
	Slope function	$Y=10.606x - 23.939$
	Regression (R ²)	0.982
96 hrs	LC ₅₀ mg/L	3.2
	NOEC mg/L	2.4
	95% Confidence Interval	2.99-3.42
	Slope function	$Y=10.97x - 13.333$
	Regression (R ²)	0.9918

LC₅₀: Lethal Concentration 50 NOEC: No Observed Effect Concentration

Conclusion:

Results obtained from the studies indicated that PAH Naphthalene is toxic to the fish, *Puntius ticto*. The results of this investigation provide long term safe levels of toxicants for fish and other aquatic life which can be estimated by using short term acute toxicity bioassays. Studies were carried out only with pure compound. Effects may increase manifold in case of derivatives of Naphthalene. However, these results may be useful for preliminary evaluation of the toxic effect of Naphthalene on the wastewaters containing this PAH. Short term bioassay results also help in calculating sub lethal doses to carry out detail bioaccumulation studies using chronic bioassays.

Acknowledgements:

1. I am immensely grateful to my mentor and guide, Late Dr. Shanta Satyanarayan for her encouragement and cooperation for carrying out this work.
2. I express thanks to my institute, St. Francis De sales Coolege, Nagpur, where the present work was carried out.

References:

1. Andelman J.B. and M.S. Suess 1970. Polynuclear aromatic hydrocarbons in the water environment. Bull. Wld. Hlth. Org, 43, 479-508.
2. Reynaud S. and Deschaux P. 2006. The effects of polycyclic aromatic hydrocarbons on the immune system of fish- A review. Aquatic Toxicology 77, No.2, 229-238.
3. Craig A.E, Mahammud Athar, Karen A.T., Dinah R and Hui Xu 1998. Susceptibility of biological effects of polyaromatic hydrocarbons is influenced by the genes of the major histocompatibility complex. Proc. Natl. Sci. USA., Vol. 95, 4015-4919.
4. Nabiha Yusuf, Laura Timares, Megan D.S., Hui Xu and Craig A.E. Acquired and innate immunity to polyaromatic hydrocarbons. Toxicol. Appl. Pharmacol.
5. Kai Yip Kong, Kwai-Chung Cheung, Chris-Kong chu Wong, Ming – Hung Wong 2005. Residues of DDTs, PAHs and some heavy metals in fish (Tilapia) collected from Hong Kong and mainland China. J-Environmental Science and Health Part A Vol.49 Issue 11, 2105-2115.
6. Ramachandran and Shahunlhata 2005. The risk of fish to exposure to polycyclic aromatic hydrocarbons from chemical dispersion of crude oil. Queens University, Kingston, Canada Page 172.
7. Kayal S. and Connell D.W. 1994. Polycyclic aromatic hydrocarbons in biota from the Brisbane River estuary, Australia. Division of Environmental Sciences, Griffith University, Nalhan, Queensland, Australia.
8. Arvo Tuvikene- 2002. Effects of chemical pollution on fish in the drainage area of the River Narva, Estonia. IJRBAT, Special Issue (2), Vol-V, July 2017 ISSN No. 2347-517X (Online) SHRI SHIVAJI SCIENCE COLLEGE, NAGPUR 916 ICRTS-2017
9. Hose, J.E., J.B. Hannah, H.W. Puffer and M.L. Landolt 1984. Histologic and skeletal abnormalities in benzo(a) pyrene treated rainbow trout alevins. Arch. Environ. Contam. Toxicol. 13, 675-684.
10. Jung – Suk lee, Kyu- Tae lee and Gyung Soo park. 2005. Acute toxicity of heavy metals, tributyltin, ammonia and polycyclic aromatic hydrocarbons to benthic amphipod *Grandidierella japonica*. Ocean Science Journal, Vol.40, No.2, 61-66.
11. S.M. Billard, Kirsten Querbach, P.V. Hodson. 2009. Toxicity of retene to early life stages of two freshwater fish species. Environmental Toxicology and Chem. Vol.18 (9), 2070-2077.
12. Couillard C.M., Laplatte B. and Pelletier E. 2009. A fish bioassay to evaluate the toxicity associated with the ingestion of benzo [a] pyrene-contaminated benthic prey. Environ. Toxicol. Chem. April; 28(4) 772-781.
13. Galtsaff P.S., Hart W.H., Patrick E.R., Strong E.R. Surles E.W., Vanitron, W.M. 1951. Bioassay methods for the evaluation of acute toxicity of industrial waste water to fish. Sewage. Ind. Waste 23, 130.

14. Sprague J.B. 1969. Review paper; Measurement of pollutant toxicity to fish: Bioassay method for acute toxicity. Water Res. 3, 793.
15. Standard methods for the examination of water and waste water (20th Edition) 1998. APHA, AWWA, WPCF, USA.
16. Lithchfield J.T. and Wilcoxon F.A.1949. A simplified method to evaluate dose effect experiments. J. Pharmacol. Exper. Thera 96, 99- 113.